


## Rabbit antibody to TrkC

<b>Code</b>	OST00179W
<b>ID Tag</b>	Rb1495-250410-WS
<b>Unit size</b>	100 ul
<b>Immunogen</b>	A synthetic peptide from the extracellular domain of rat TrkC conjugated to blue carrier protein was used as the antigen. The peptide is homologous in mouse and human.
<b>Conjugate</b>	Unconjugated antibody
<b>Also known</b>	Neurotrophic tyrosine kinase receptor type 3, TrkC tyrosine kinase, GP145-TrkC, Trk-C, NT-3 growth factor receptor, NTRK3
<b>Host</b>	NZ white rabbit
<b>Purity</b>	Whole serum
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Polyclonal, whole serum
<b>Applications</b>	IHC. A dilution of 1 : 1000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.
<b>Specificity</b>	Specific for TrkC.
<b>Spcs X-react.</b>	Rat, mouse, human. Other species not yet tested.
<b>Format</b>	Lyophilised
<b>Reconstitution</b>	Reconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.
<b>Storage</b>	Maintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and refrigerated at 2-8C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles.
<b>Expiry Date</b>	12 months after reconstitution
<b>Shipping</b>	This item will be shipped to you at ambient temperature in a lyophilised form.
<b>Limitation</b>	For research use only



IF on rat DRG at 1 : 1000 dilution using Rabbit antibody to extracellular domain of TrkC (OST00179W), DAPI counterstained.

---

### Related Products

OST00293G	Rabbit antibody to TrkC
OST00119W	Sheep antibody to TrkC
OST00120G	Sheep antibody to TrkC