


Rabbit antibody to AQP3

Code	OSA00087W
ID Tag	Rb968-280609-WS
Unit size	100 µl
Immunogen	A synthetic peptide from the c-terminal region of rat Aquaporin 3 (AQP3) conjugated to an immunogenic carrier protein was used as the antigen.
Conjugate	Unconjugated antibody
Also known	Aquaporin-3, AQP-3, AQP3, Aquaporin 3, Aquaporin3
Host	NZ white rabbit
Purity	Whole serum
Clonality	Polyclonal
Isotype	Polyclonal, whole serum
Applications	IHC, WB. A dilution of 1 : 300 to 1 : 2000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.
Specificity	Specific for AQP3.
Spcs X-react.	Rat, mouse. Other species not yet tested.
Format	Lyophilised
Reconstitution	Reconstitute in 100 µl of sterile water. Centrifuge to remove any insoluble material.
Storage	Maintain the lyophilised/reconstituted antibodies frozen at -20°C for long term storage and refrigerated at 2-8°C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles.
Expiry Date	12 months after reconstitution
Shipping	This item will be shipped to you at ambient temperature in a lyophilised form.
Limitation	For research use only



```

/* Font Definitions */
@font-face
{font-family:Arial;
panose-1:2 11 6 4 2 2 2 2 4;
mso-font-charset:0;
mso-generic-font-family:auto;
mso-font-pitch:variable;
mso-font-signature:3 0 0 0 1 0;}
@font-face
{font-family:Cambria;
panose-1:2 4 5 3 5 4 6 3 2 4;
mso-font-charset:0;
mso-generic-font-family:auto;
mso-font-pitch:variable;
mso-font-signature:3 0 0 0 1 0;}
/* Style Definitions */
p.MsoNormal, li.MsoNormal, div.MsoNormal

```

Related Products

OSA00140W Goat antibody to AQP3
margin-right:0cm;
margin-bottom:10.0pt;
margin-left:0cm;

Rabbit antibody to AQP3

```
mso-pagination:widow-orphan;  
font-size:12.0pt;  
font-family:"Times New Roman";  
mso-ascii-font-family:Arial;  
mso-fareast-font-family:Cambria;  
mso-fareast-theme-font:minor-latin;  
mso-hansi-font-family:Arial;  
mso-bidi-font-family:"Times New Roman";  
mso-bidi-theme-font:minor-bidi;}  
@page Section1  
{size:612.0pt 792.0pt;  
margin:72.0pt 90.0pt 72.0pt 90.0pt;  
mso-header-margin:36.0pt;  
mso-footer-margin:36.0pt;  
mso-paper-source:0;}  
div.Section1  
{page:Section1;}  
-->
```

IHC on paraffin sections of rat kidney tissue using Rabbit antibody to AQP3: OSA00087W.
HIER: 1 mM EDTA, pH 8 for 20 min using Thermo PT Module.
Blocking: 0.2% LFDM in TBST filtered thru 0.2 μ m.
Detection was done using Novolink HRP polymer from Leica following manufacturer's instructions.
Primary antibody: dilution 1: 1000, incubated 30 min at RT (using Autostainer).
Sections were counterstained with Harris Hematoxylin.

Related Products

OSA00140W Goat antibody to AQP3