


## Rabbit antibody to 5HT5A-R

<b>Code</b>	OSS00198W
<b>ID Tag</b>	Rb1153-260909-WS
<b>Unit size</b>	100 ul
<b>Immunogen</b>	A synthetic peptide from n-terminal region of human 5HT5A receptor (Serotonin receptor 5A) conjugated to blue carrier protein was used as the antigen.
<b>Conjugate</b>	Unconjugated antibody
<b>Also known</b>	5-hydroxytryptamine receptor 5A, 5-HT5A, Serotonin receptor 5A, HTR5A
<b>Host</b>	NZ white rabbit
<b>Purity</b>	Whole serum
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Polyclonal, whole serum
<b>Applications</b>	IHC, WB. A dilution of 1: 300 to 1: 2000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.
<b>Specificity</b>	Specific for 5HT5A receptor.
<b>Spcs X-react.</b>	Human, rat, mouse. Other species not yet tested.
<b>Format</b>	Lyophilised. This product contains 0.02% benzalkonium chloride.
<b>Reconstitution</b>	Reconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.
<b>Storage</b>	Maintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and refrigerated at 2-8C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles.
<b>Expiry Date</b>	12 months after reconstitution
<b>Shipping</b>	This item will be shipped to you at ambient temperature in a lyophilised form.
<b>Limitation</b>	For research use only



IHC on paraffin sections of human uterus.  
 HIER: 1 mM EDTA, pH 8 for 20 min using Thermo PT Module.  
 Blocking: 0.2% LFDM in TBST filtered thru 0.2 µm.  
 Detection was done using Novolink HRP polymer from Leica following manufacturers instructions.  
 Primary antibody: dilution 1: 1000, incubated 30 min at RT (using Autostainer).  
 Sections were counterstained with Harris Hematoxylin.