


Rabbit antibody to HPCA

| | |
|-----------------------|---|
| Code | OSH00033W |
| ID Tag | Rb1215-051209-WS |
| Unit size | 100 ul |
| Immunogen | A synthetic peptide from the c-terminal region of human Hippocalcin (HPCA) conjugated to blue carrier protein was used as the antigen. |
| Conjugate | Unconjugated antibody |
| Also known | HPCA, Neuron-specific calcium-binding protein hippocalcin, Calcium-binding protein BDR-2, BDR2 |
| Host | NZ white rabbit |
| Purity | Whole serum |
| Clonality | Polyclonal |
| Isotype | Polyclonal, whole serum |
| Applications | IHC, WB. A dilution of 1: 200 to 1: 2000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications. |
| Specificity | Specific for hippocalcin. |
| Spcs X-react. | Human, rat, mouse. Other species not yet tested. |
| Format | Lyophilised |
| Reconstitution | Reconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material. |
| Storage | Maintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and refrigerated at 2-8C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles. |
| Expiry Date | 12 months after reconstitution |
| Shipping | This item will be shipped to you at ambient temperature in a lyophilised form. |
| Limitation | For research use only |



IHC on rat spinal cord (paraffin section) using Rabbit antibody to HPCA (OSH00033W) at 1 : 200 dilution incubated overnight at 4C, developed with DAB/Ni and counterstained with hematoxylin.