

Rabbit antibody to RYR1 (1360-1400)

Code OSR00366W

ID Tag Rb1666-211110-WS

Unit size 100 μl

Immunogen A synthetic peptide from aa region 1360-1400 of human RYR1 conjugated to blue carrier protein

was used as the antigen. The peptide is homologous in rat and mouse.

Conjugate Unconjugated antibody

Also known RYR-1, RyR1, Skeletal muscle calcium release channel, Skeletal muscle-type ryanodine receptor,

RYDR

Host NZ white rabbit
Purity Whole serum
Clonality Polyclonal

Isotype Polyclonal, whole serum

Applications IHC, WB (confirmed by recombinant protein). A dilution of 1:1000 to 1:2000 is recommended.

The optimal dilution should be determined by the end user. Not yet tested in other applications.

Specificity Specific for RYR1.

Spcs X-react. Human, mouse, rat. Other species not yet tested.

Format Lyophilised

Reconstitution Reconstitute in 100 µl of sterile water. Centrifuge to remove any insoluble material.

Storage Maintain the lyophilised/reconstituted antibodies frozen at -20°C for long term storage and


refrigerated at 2-8°C for a shorter term. When reconstituting, glycerol (1:1) may be added for an

additional stability. Avoid freeze and thaw cycles.

Expiry Date 12 months after reconstitution

Shipping This item will be shipped to you at ambient temperature in a lyophilised form.

Limitation For research use only



IHC-P on paraffin sections of rat DRG.

The animal was perfused using Autoperfuser at a pressure of 130 mmHg with 300 ml 4% FA before being processed for paraffin embedding. HIER: Tris-EDTA, pH 9 for 20 min using Thermo PT Module.

Blocking: 0.2% LFDM in TBST filtered thru 0.2 $\mu m.$

Detection was done using Novolink HRP polymer from Leica following manufacturers instructions; DAB chromogen: Candela DAB chromogen from Osenses.

Primary antibody: dilution 1: 1000, incubated 30 min at RT using Autostainer.

Sections were counterstained with Harris Hematoxylin.