


Rabbit antibody to TAAR2

| | |
|-----------------------|---|
| Code | OSR00080W |
| ID Tag | Rb1398-250410-WS |
| Unit size | 100 µl |
| Immunogen | A synthetic peptide from n-terminal region of human trace amine-associated receptor 2 (TAAR2) conjugated to an immunogenic carrier protein was used as the antigen. |
| Conjugate | Unconjugated antibody |
| Also known | Trace amine-associated receptor 2, TAAR2, GPR58 |
| Host | NZ white rabbit |
| Purity | Whole serum |
| Clonality | Polyclonal |
| Isotype | Polyclonal, whole serum |
| Applications | IHC, WB. A dilution of 1 : 300 to 1 : 2000 is recommended. The optimal dilution should be determined by the end user. |
| Specificity | Specific for TAAR2. |
| Spcs X-react. | Human. Other species not yet tested. |
| Format | Lyophilised |
| Reconstitution | Reconstitute in 100 µl of sterile water. Centrifuge to remove any insoluble material. |
| Storage | Maintain the lyophilised/reconstituted antibodies frozen at -20°C for long term storage and refrigerated at 2-8°C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles. |
| Expiry Date | 12 months after reconstitution |
| Shipping | This item will be shipped to you at ambient temperature in a lyophilised form. |
| Limitation | For research use only |



IHC on paraffin sections of human brain tissue. HIER: Tris-EDTA, pH 9 for 20 min using Thermo PT Module. Blocking: 0.2% LFDM in TBST filtered thru 0.2 µm. Detection was done using Novolink HRP polymer from Leica following manufacturers instructions. Primary antibody: dilution 1: 1000, incubated 30 min at RT (using Autostainer). Sections were counterstained with Harris Hematoxylin.