

Sheep antibody to TRPM4

Code OST00033W

ID Tag Sh238-1-280310-WS

Unit size 100 μl

Immunogen A synthetic peptide from human TRPM4 conjugated to blue carrier protein was used as the

antigen.

Conjugate Unconjugated antibody

Also known Transient receptor potential cation channel subfamily M member 4, long transient receptor

potential channel 4, hTRPM4, melastatin-4, calcium-activated non-selective cation channel 1,

TRPM4, TRPM4B, FLJ20041

Host Sheep

Purity Whole serum Clonality Polyclonal

Isotype Polyclonal, whole serum

Applications IHC, WB. A dilution of 1: 300 to 1: 2000 is recommended. The optimal dilution should be

determined by the end user. Not yet tested in other applications.

Specificity Specific for TRPM4.

Spcs X-react. Human. Other species not yet tested.

Format Lyophilised

Reconstitution Reconstitute in 100 µl of sterile water. Centrifuge to remove any insoluble material.

Storage Maintain the lyophilised/reconstituted antibodies frozen at -20°C for long term storage and


refrigerated at 2-8°C for a shorter term. When reconstituting, glycerol (1:1) may be added for an

additional stability. Avoid freeze and thaw cycles.

Expiry Date 12 months after reconstitution

Shipping This item will be shipped to you at ambient temperature in a lyophilised form.

Limitation For research use only



A cell cytospin of human lung cancer cell line A549 was fixed with cold acetone for 90 seconds and air-dried. Cells were incubated with the blocking buffer (PBS containing 5% FCS) for 30 minutes at room temperature. Cells were then washed once in PBS and incubated with primary antibody, diluted 1:100 in the blocking buffer, for 30 minutes. Slides were washed 3X in PBS and incubated with Goat anti-sheep conjugated to Alexa-488, diluted 1:200 in blocking buffer, for 30 minutes at room temperature in the dark. Slides were washed as above and mounting media (10% Glycerol in PBS) containing Hoechst 33258 1 μ g/ml was used for nuclear counterstaining. Fluorescent cell staining were analysed using a Olympus microscope and the analysis LS Research software.