


## Rabbit antibody to VACHT

|                       |                                                                                                                                                                                                                                                 |
|-----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Code</b>           | OSV00001W                                                                                                                                                                                                                                       |
| <b>ID Tag</b>         | Rb1170-260909-WS                                                                                                                                                                                                                                |
| <b>Unit size</b>      | 100 ul                                                                                                                                                                                                                                          |
| <b>Immunogen</b>      | A synthetic peptide from human VACHT conjugated to blue carrier protein was used as the antigen. The antigen is homologous in rat and mouse.                                                                                                    |
| <b>Conjugate</b>      | Unconjugated antibody                                                                                                                                                                                                                           |
| <b>Also known</b>     | Vesicular acetylcholine transporter, solute carrier family 18 member 3, SLC18A3, MGC12716                                                                                                                                                       |
| <b>Host</b>           | NZ white rabbit                                                                                                                                                                                                                                 |
| <b>Purity</b>         | Whole serum                                                                                                                                                                                                                                     |
| <b>Clonality</b>      | Polyclonal                                                                                                                                                                                                                                      |
| <b>Isotype</b>        | Polyclonal, whole serum                                                                                                                                                                                                                         |
| <b>Applications</b>   | IHC, WB. A dilution of 1: 2000 is recommended for WB and to 1:1000 for IHC-P. The optimal dilution should be determined by the end user. Not yet tested in other applications.                                                                  |
| <b>Specificity</b>    | Specific for VACHT.                                                                                                                                                                                                                             |
| <b>Spcs X-react.</b>  | Human, mouse, rat, marmoset. Other species not yet tested.                                                                                                                                                                                      |
| <b>Format</b>         | Lyophilised                                                                                                                                                                                                                                     |
| <b>Reconstitution</b> | Reconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.                                                                                                                                                           |
| <b>Storage</b>        | Maintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and refrigerated at 2-8C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles. |
| <b>Expiry Date</b>    | 12 months after reconstitution                                                                                                                                                                                                                  |
| <b>Shipping</b>       | This item will be shipped to you at ambient temperature in a lyophilised form.                                                                                                                                                                  |
| <b>Limitation</b>     | For research use only                                                                                                                                                                                                                           |



IHC-P on paraffin sections of rat spinal cord.

The animal was perfused using Autoperfuser at a pressure of 130 mmHg with 300 ml 4% FA before being processed for paraffin embedding. HIER: Tris-EDTA, pH 9 for 20 min using Thermo PT Module.

Blocking: 0.2% LFDM in TBST filtered thru 0.2 µm.

Detection was done using Novolink HRP polymer from Leica following manufacturers instructions; DAB chromogen: Candela DAB chromogen from Osenses.

Primary antibody: dilution 1: 1000, incubated 30 min at RT using Autostainer.

Sections were counterstained with Harris Hematoxylin.